

Lop/r	80/20 mg/ml	2 ml + 1 ml	2 ml + 1 ml	2 ml + 2 ml	3 ml + 2 ml	3 mL+ 3 ml	200/50 (adult)	2 + 1	2 + 1
-------	-------------	-------------	-------------	-------------	-------------	------------	----------------	-------	-------

Note: Higher doses of Lop/r may be required when co-administered with enzyme-inducing medicines such as NVP, EFV; n/r = not recommended.

Medicine	Strength of paediatric tablet (mg)	Number of tablets by weight band once daily Children 3 years and over					Strength of paediatric tablet (mg)	Number of tablets by weight band once daily (PM dosing preferred)	
		n/r	n/r	1x200 mg	1x200 mg +	1x200 mg +		1.5x200 mg +	2x200 mg
EFV	200, 50	n/r	n/r	1x200 mg	1x200 mg +	1x200 mg +	200, 50	1.5x200 mg +	2x200 mg
					1x50 mg	2x50 mg		1x50 mg	
EFV	200, 50	n/r	n/r	0 + 1	0 + 2	0 + 3		0 + 2.5	0 + 2

(Note: Pediatric Dosing Chart on preceding page was adapted from: International Center for AIDS Care and Treatment Programs, Global AIDS Program, Baylor International Pediatric AIDS Initiative, Pediatric Antiretroviral Dosing in Resource-Limited Settings, Updated November 2006. Available at: http://www.cdc.gov/globalAIDS/pa_pmtct_pediatric.htm.)

